



King's Research Portal

DOI:

[10.1016/j.bbi.2017.05.006](https://doi.org/10.1016/j.bbi.2017.05.006)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Borsini, A., Alboni, S., Tojo, L. M., Horowitz, M. A., Su, K-P., Cannazza, G., Pariente, C. M., & Zunszain, P. A. (2017). Rescue of IL-1induced reduction of human neurogenesis by omega-3 fatty acids and antidepressants. *Brain, Behavior, and Immunity*, 65, 230-238. <https://doi.org/10.1016/j.bbi.2017.05.006>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Full-length Article

Rescue of IL-1 β -induced reduction of human neurogenesis by omega-3 fatty acids and antidepressantsAlessandra Borsini^a, Silvia Alboni^b, Mark A. Horowitz^a, Luis M. Tojo^a, Giuseppe Cannazza^b, Kuan-Pin Su^c, Carmine M. Pariante^{a,d}, Patricia A. Zunszain^{a,*}^a Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK^b Department of Life Sciences, University of Modena and Reggio Emilia, Modena, Italy^c Department of Psychiatry & Mind-Body Interface Laboratory (MBI-Lab), China Medical University Hospital, College of Medicine, China Medical University, Taichung, Taiwan^d South London and Maudsley NHS Foundation Trust, Denmark Hill, Camberwell, London, UK

ARTICLE INFO

Article history:

Received 10 January 2017

Received in revised form 24 April 2017

Accepted 8 May 2017

Available online 18 May 2017

Keywords:

Fish oil

Cytokines

PUFA

Immune

IL-1 beta

Sertraline

Venlafaxine

Neurogenic

Kynurenine-pathway

ABSTRACT

Both increased inflammation and reduced neurogenesis have been associated with the pathophysiology of major depression. We have previously described how interleukin-1 (IL-1) β , a pro-inflammatory cytokine increased in depressed patients, decreases neurogenesis in human hippocampal progenitor cells. Here, using the same human *in vitro* model, we show how omega-3 (ω -3) polyunsaturated fatty acids and conventional antidepressants reverse this reduction in neurogenesis, while differentially affecting the kynurenine pathway. We allowed neural cells to proliferate for 3 days and further differentiate for 7 days in the presence of IL-1 β (10 ng/ml) and either the selective serotonin reuptake inhibitor sertraline (1 μ M), the serotonin and norepinephrine reuptake inhibitor venlafaxine (1 μ M), or the ω -3 fatty acids eicosapentaenoic acid (EPA, 10 μ M) or docosahexaenoic acid (DHA, 10 μ M). Co-incubation with each of these compounds reversed the IL-1 β -induced reduction in neurogenesis (DCX- and MAP2-positive neurons), indicative of a protective effect. Moreover, EPA and DHA also reversed the IL-1 β -induced increase in kynurenine, as well as mRNA levels of indolamine-2,3-dioxygenase (IDO); while DHA and sertraline reverted the IL-1 β -induced increase in quinolinic acid and mRNA levels of kynurenine 3-monooxygenase (KMO). Our results show common effects of monoaminergic antidepressants and ω -3 fatty acids on the reduction of neurogenesis caused by IL-1 β , but acting through both common and different kynurenine pathway-related mechanisms. Further characterization of their individual properties will be of benefit towards improving a future personalized medicine approach.

Crown Copyright © 2017 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Hippocampal neurogenesis has been widely associated with depression: reduced levels have been reported in depressed patients as well as in animal models of depression (Boldrini et al., 2009; Kempermann et al., 2008), and increasing its levels appears sufficient to reduce depressive-like behaviours (Hill et al., 2015). Several factors can regulate neurogenesis: it can be stimulated by antidepressant drugs (Warner-Schmidt and Duman, 2006; Sahay and Hen, 2007; Malberg and Blendy, 2005) and negatively affected by pro-inflammatory cytokines such as interleukin-1 (IL-1) β (Goshen et al., 2008; Koo and Duman, 2008; Kuzumaki et al., 2010). This cytokine is of importance, as increased levels have been shown in peripheral blood and cere-

brospinal fluid of depressed patients (Capuron and Miller, 2004; Howren et al., 2009; Maes et al., 2009; Miller et al., 2009; Mossner et al., 2007; Raison et al., 2006; Schiepers et al., 2005). Notably, higher levels of IL-1 β are known to affect synaptic transmission, causing impairments of learning and memory (Yirmiya and Goshen, 2011).

One of the underlying mechanisms by which IL-1 β and other pro-inflammatory cytokines can affect neurogenesis is through alterations of the kynurenine pathway (see Fig. 1 for a simplified description). This pathway starts from the essential amino acid tryptophan and can generate the neurotoxic quinolinic acid (QUIN), a putative NMDA receptor antagonist (Schwarcz et al., 2012), whose accumulation can be prevented by production of picolinic acid (PIC), a potentially neuroprotective metabolite. Dysregulation in levels of several kynurenine pathway metabolites has been described in depression and other psychiatric disorders. For example, increased levels of QUIN have been detected in animal

* Corresponding author.

E-mail address: patricia.zunszain@kcl.ac.uk (P.A. Zunszain).




Fig. 1. Simplified kynurenine pathway of tryptophan metabolism. ACMSD: aminocarboxymuconate semialdehyde decarboxylase, IDO: indolamine-2,3-dioxygenase, KAT: kynurenine aminotransferase, KMO: kynurenine 3-monooxygenase or kynurenine 3-hydroxylase, KYNU: kynureninase.

models of depressive-like behavior (Fischer et al., 2015), in hepatitis C patients with depression induced by IFN- α (Raison et al., 2010) and in the cerebrospinal fluid (CSF) of suicide attempters (Erhardt et al., 2013), who have also been shown to have reduced PIC and a decreased PIC/QUIN ratio in both CSF and blood (Brundin et al., 2016). Similarly, increased levels of kynurenine have been detected in depression induced by IFN- α (Capuron et al., 2003) and in the hippocampus of patients with Alzheimer's disease (Guillemin et al., 2005), while the peripheral ratio of kynurenic acid/QUIN ratio has been shown to be reduced in depressed individuals (Meier et al., 2016; Savitz et al., 2015).

Considering the regulatory effects of these metabolites, the role of enzymes of the pathway that control their production is of significant interest. For example, inhibition of IDO, the first enzyme of the pathway, reduced the depression-like behavior induced by inflammation via peripheral (O'Connor et al., 2009) or intracerebral (Dobos et al., 2012) administration of lipopolysaccharide (LPS) in mice. Furthermore, inhibition of KMO reversed cognitive and motor deficits measured in neurodegeneration, using mouse models of Alzheimer's disease and Huntington's disease (Zwilling et al., 2011). Interestingly, KMO-deficient mice have been shown to be protected from inflammation-induced depressive-like behavior (Heisler and O'Connor, 2015; Parrott et al., 2016).

Our previous research has shown that IL-1 β decreases neurogenesis while increasing levels of expression of all neurotoxicity-associated enzymes of the kynurenine pathway. This effect was partially reversed by KMO inhibition, suggesting that activation of the kynurenine pathway is indeed contributing to this cytokine-induced reduction of neurogenesis (Zunszain et al., 2012). Using the same *in vitro* model, we have also described how reduced neurogenesis induced by glucocorticoids is reversed by the SSRI sertraline (Anacker et al., 2011). Considering that IL-1 β levels decrease in patients upon antidepressant treatment (Hannestad et al., 2011), we here explore the possibility that a regulation of neurogenesis in inflammatory conditions may be a common underlying pathway of antidepressant action. We investigate two antidepressants, the SSRI sertraline and the SNRI venlafaxine, as well as the two principal ω -3 polyunsaturated fatty acids components of fish oils, eicosapentaenoic acid (EPA) and docosahexanoic acid (DHA), which have been associated with the pathogenesis and treatment of depression (Su et al., 2014; Su, 2012, 2009; Lin et al., 2012, 2010; Lu et al., 2010). We have recently

shown that these four compounds have different immunomodulating properties upon co-treatment with IL-1 β : while all were associated with a corresponding decrease in the NF- κ B activation caused by IL-1 β alone, venlafaxine and EPA showed mostly anti-inflammatory properties but sertraline and DHA displayed mostly pro-inflammatory properties (Horowitz et al., 2015). In addition to looking at their effects on neurogenesis in the presence of an inflammatory insult, we here extend the study of their properties, evaluating additionally whether they regulate the metabolites and enzymes of the kynurenine pathway.

2. Materials and methods

2.1. Cell culture

All experiments were performed with the multipotent human hippocampal progenitor cell line HPC0A07/03C provided by ReNeuron (Surrey, UK). Cells were grown in reduced modified media consisting of Dulbecco's Modified Eagle's Media/F12 (DMEM:F12, Invitrogen, Paisley, UK) supplemented with 0.03% human albumin (Baxter Healthcare, Compton, UK), 100 μ g/mL human apo-transferrin, 16.2 μ g/mL human putrescine diHCl, 5 μ g/mL human recombinant insulin, 60 ng/mL progesterone, 2 mM L-glutamine and 40 ng/mL sodium selenite. To maintain proliferation, 10 ng/mL human basic fibroblast growth factor (bFGF), 20 ng/mL human epidermal growth factor (EGF) and 100 nM 4-hydroxytamoxifen (4-OHT) were added. Cells were grown in 75 cm² filtered cap culture flasks (Nunc, Roskilde, Denmark) at 37 °C in 5% CO₂ and regularly passaged at 80% confluence, until being transferred to plates for the differentiation experiments.

2.2. Assays with antidepressants and ω -3 fatty acids

To assess whether antidepressants and ω -3 fatty acids could abrogate changes in neuronal differentiation caused by IL-1 β exposure, HPC0A07/03C cells were plated on 96 well plates (Nunc) at a density of 1000 cells per well. Cells were cultured in the presence of EGF, bFGF and 4-OHT and treated with IL-1 β (10 ng/mL) either alone or in combination with the antidepressant venlafaxine (1 μ M) or sertraline (1 μ M), or one of the ω -3 fatty acids, EPA (10 μ M) or DHA (10 μ M). Antidepressants and ω -3 fatty acids were dissolved in EtOH. All treatments had the same vehicle, including

0.1% ethanol, to exclude the possibility of any differences observed being the consequence of differing concentrations of solvents. Treatment conditions and doses were chosen as described in our previous work (Horowitz et al., 2015). After 3 days, cell-conditioned media was collected and stored at -80°C for subsequent measurement of kynurenine metabolites, cells were washed and treatment with the cytokine and each of the four compounds was repeated in media without growth factors and 4-OHT for 7 subsequent days. At the end of the total incubation time (10 days), cells were fixed with 4% PFA to be further processed by immunocytochemistry. Treatment with the cytokine and all drugs was also conducted in 6 well plates to extract RNA after 24 h in proliferation media, in order to investigate changes in gene expression.

2.3. Immunocytochemistry

Neuronal differentiation and maturation were assessed by doublecortin (DCX) and microtubulin-associated protein-2 (MAP2) immunocytochemistry. Briefly, PFA-fixed cells were incubated in blocking solution (5% normal donkey serum, Scientific Laboratory Supplies Ltd, Nottingham, UK) in PBS containing 0.3% Triton-X for 2 h at room temperature, and with primary antibodies [rabbit anti-DCX, 1:500; mouse anti-MAP2 (HM), 1:500, Abcam, Cambridge, UK] at 4°C overnight. Specificity of the MAP2 (HM) antibody for mature neurons in our cell culture was previously confirmed (Anacker et al., 2011). Cells were incubated sequentially in blocking solution for 30 min, secondary antibodies (Alexa 488 donkey anti-rabbit; 1:1000; Alexa donkey 555 anti-mouse, 1:1000, Invitrogen) for 1 h, and Hoechst 33342 dye (0.01 mg/ml, Invitrogen) for 5 min at room temperature. The number of DCX- and MAP2-positive cells over total Hoechst 33342 positive cells was counted with an automated approach using CellInsight NXT High Content Screening (HCS) Platform (ThermoScientific). Negative controls were incubated with unspecific mouse IgGs (1:500, control for MAP2), rabbit IgGs (1:500, control for DCX) in place of the specific primary antibody.

2.4. RNA isolation and cDNA synthesis

RNA was isolated using the RNeasy Micro Kit (Qiagen, Crawley, UK) following the manufacturer's instructions and samples were kept frozen at -80°C until further use. RNA quantity and quality were assessed by evaluation of the A260/280 and A260/230 ratios using a Nanodrop spectrometer (NanoDrop Technologies, Wilmington, DE, USA). Superscript III enzyme (Invitrogen, Carlsbad, CA, USA) was used to reverse transcribe 1 μg total RNA as previously described (Anacker et al., 2011).

2.5. Quantitative Real Time PCR (qPCR) analyses

qPCR was performed using Predesigned TaqMan Gene Expression Assay probes (Life Technologies) with TaqMan Universal PCR Master Mix with UNG (Life Technologies), using Chromo 4 DNA instrument from BioRad. The expression of target genes indoleamine 2,3-dioxygenase (IDO), kynurenine 3-monooxygenase (KMO), kynureninase (KYNU) and aminocarboxymuconate semialdehyde decarboxylase (ACMSD) was normalized to the expression levels of beta actin (ACTB) and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) as references. The relative expression levels of target genes detected were calculated using the Pfaffl method (Bustin et al., 2005), with data normalized to the geometric mean of the housekeeping genes and expressed as fold change compared with the control sample. Three independent experiments were conducted on independent cultures, and each sample was tested in duplicate.

2.6. Sample preparation for metabolite analysis

After 3 days of proliferation, before media was changed to allow differentiation for 7 subsequent days, the cell media from each experimental condition was removed and centrifuged ($2000 \times \text{rpm}$, 5 min) to pellet and remove dead cells. This supernatant was then assayed for metabolites of the kynurenine pathway, by adding an equal volume of ice cold 1 M perchloric acid (HClO_4) fortified with deuterated tryptophan as internal standard (final concentration 1 μM); samples were centrifuged ($15,000 \times \text{g}$, 10 min at 4°C) and the supernatants were collected for direct injection into the LC-MS/MS.

2.7. Liquid chromatography

The analyses of tryptophan (Trp), kynurenine (KYN), anthranilic acid (ANA), picolinic acid (PIC) and quinolinic acid (QUIN) were performed using an Agilent HP 1200 liquid chromatograph (Agilent, Milan, Italy) consisting of a binary pump, an autosampler and a thermostated column compartment. Chromatographic separations were carried out using a Discovery HS-F5 column ($150 \times 4.6 \text{ mm}$, 5 μm , Supelco, Milan, Italy) using 0.1% formic acid in water and acetonitrile (ACN) as mobile phase. The HPLC analyses were carried out using a linear elution profile of 15 min from 5% to 90% of ACN. The column was washed with 90% ACN for 3.5 min, then equilibrated for 5 min with 5% ACN. The flow rate was 0.5 mL/min. The injection volume was 40 μL . An Agilent 6410 triple quadrupole-mass spectrometer with an electrospray ion source operating in positive mode was used for detection. The optimized source parameters for MS analysis were: drying gas temperature 350°C and gas flow 12 L/min, nebulizer gas flow pressure 35 psi and capillary voltage 4500 V. The optimized fragmentor voltage was 66 V and the collision energies were 4, 4, 8, 8, 8, 24 and 8 eV for Trp, KYN, ANA, PIC, QUIN and internal standard, respectively. The SRM pairs were 205 \rightarrow 188 (118), 209 \rightarrow 192 (94), 138 \rightarrow 120 (65), 124 \rightarrow 106 (51) and 168 \rightarrow 78 (114) and 210 \rightarrow 192 for Trp, KYN, ANA, PIC, QUIN and internal standard, respectively. The calibration curves were constructed using seven calibration standards and were linear over the concentration range of 0.001–2.000 μM for ANA, 0.01–5.00 μM for KYN, PIC and QUIN and 0.1–20 μM for TRP, with a correlation coefficient (r^2) ≥ 0.9983 for all analytes. The limit of detection at a signal-to-noise ratio of 3:1 and lower limit of quantitation were 0.0005 and 0.0010 μM for ANA; 0.005 and 0.010 μM for KYN, PIC and QUIN; 0.05 and 0.1 μM for TRP.

2.8. Drugs and reagents

All drugs and reagents were purchased from Sigma–Aldrich (St Louis, MO, USA), unless otherwise stated. IL-1 β , EGF and bFGF were purchased from Peprotech (London, UK).

2.9. Statistical analysis

Data are presented as mean \pm SEM. All statistical analyses were performed with GraphPad Prism 7 on four independent biological replicates. One-Way ANOVA with Bonferroni's post hoc test was used for multiple comparisons among treatment groups. Student's *t*-test was used to compare means of two independent treatment groups; *p*-values of <0.05 were considered significant.

3. Results

3.1. Antidepressants and ω -3 fatty acids prevent the IL-1 β -induced reduction of neurogenesis

Treatment of cells with IL-1 β (10 ng/mL) for three days of proliferation followed by seven days of differentiation reduced the percentage of DCX-positive and MAP2-positive cells by 28% and 40%, respectively (Fig. 2a). These changes were similar to those we have previously reported (Zunszain et al., 2012). To establish whether the ω -3 fatty acids EPA and DHA, and the monoaminergic antidepressants venlafaxine and sertraline, were able to reverse these detrimental effects of IL-1 β cells were co-incubated with the cytokine and each of the compounds at doses previously established (Horowitz et al., 2015). Co-incubation of cells with IL-1 β and either EPA (10 μ M), DHA (10 μ M), sertraline (1 μ M) or venlafaxine (1 μ M) reversed the decrease in the percentage of DCX-positive neurons caused by IL-1 β to +50%, +70%, +53% and +53%, respectively (Fig. 2b). Of note, individual treatment with either ω -3 fatty acids or antidepressants did not affect the number of DCX-positive cells (data not shown). Similarly, the reduction of MAP2-positive cells caused by IL-1 β was reversed upon co-incubation with all drugs to +94% with EPA, +88% with DHA, +79% with sertraline and +67% with venlafaxine (Fig. 2c). Treatment with ω -3 fatty acids alone did not affect the number of MAP2-positive cells. In accor-

dance with our previous data (Anacker et al., 2011), sertraline significantly increased the number of MAP2-positive cells by +7%, when compared with control (data not shown). Venlafaxine also enhanced MAP2-positive staining by +8% (data not shown). Although antidepressants alone were able to affect the number of MAP2-positive cells, this effect was much lower than what we report upon co-incubation with IL-1 β and both antidepressants (+79% for sertraline and +67% for venlafaxine). Therefore, these findings are consistent with the notion that antidepressant compounds may play an important role in reversing the negative effects of inflammation on neurogenesis.

3.2. IL-1 β treatment modulates levels of metabolites and enzymes of the kynurenine pathway

Treatment of cells with IL-1 β affected the production of metabolites and enzymes of the kynurenine pathway. Specifically, IL-1 β increased kynurenine levels detected in the supernatant of proliferating cells exposed to the cytokine for 3 days by 96%, and increased levels of quinolinic acid by 208%, compared with vehicle-treated cells, and decreased levels of anthranilic acid by 28%. No significant changes were observed for either tryptophan or picolinic acid in the conditioned media (Fig. 3a). Treatment with IL-1 β increased levels of expression of enzymes involved in the production of quinolinic acid, namely IDO, KMO and KYNU, with




Fig. 2. Monoaminergic antidepressants and ω -3 fatty acids reverse the IL- β -induced reduction of human hippocampal cells neurogenesis. Treatment with IL-1 β decrease DCX-positive cells (immature neuronal phenotype), and MAP2 positive cells (mature neuronal phenotype) (a). Treatment with EPA, DHA, sertraline or venlafaxine reverse the IL-1 β -induced decrease in DCX positive cells (b), and MAP2-positive cells (c); **** $p < 0.0001$.




Fig. 3. IL-1 β modulates the production of metabolites and enzymes of the kynurenine pathway. Treatment with IL-1 β affected the kynurenine pathway by (a) increasing levels of kynurenine and quinolinic acid and decreasing levels of anthranilic acid released into the supernatant, and (b) up-regulating the expression of IDO, KMO, KYNU and ACMSD; * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$.

fold changes of 15, 13 and 9, respectively, in line with what we have shown before (Zunszain et al., 2012). We here extended our evaluation of enzymes measuring ACMSD, which is responsible for limiting the formation of quinolinic acid by production of the neuroprotective metabolite picolinic acid. Treatment with the cytokine caused a 3-fold increase in levels of expression of this enzyme (Fig. 3b).

3.3. Antidepressants and ω -3 fatty acids differentially modulate levels of metabolites of the kynurenine pathway

To investigate whether the ability of antidepressants and ω -3 fatty acids to counteract the negative effects of IL-1 β on neurogenesis involved effects on the kynurenine pathway, we first measured their action on metabolites. No compound significantly modulated the amount of tryptophan, highly present in the culture media, with respect to IL-1 β exposed cells (data not shown), however the increase in levels of kynurenine caused by the cytokine was partially reverted by co-incubation with both ω -3 fatty acids (from 96% with IL-1 β alone to 47% when co-incubated with EPA and 41% with DHA) (Fig. 4a). On the contrary, the decrease in anthranilic acid was abrogated by co-treatment with sertraline (from -28% to +12%) (Fig. 4b). Similarly, DHA and sertraline partially abrogated the rise in quinolinic acid caused by the cytokine (from +208% with IL-1 β alone to +45% when co-incubated with DHA and +49% with sertraline), whereas EPA and venlafaxine showed a non-significant reduction in quinolinic acid levels (Fig. 4c). There was no significant effect of either ω -3 fatty acids or antidepressants on levels of picolinic acid (Fig. 4d).

3.4. Antidepressants and ω -3 fatty acids differentially regulate enzymes of the kynurenine pathway

Both antidepressants and ω -3 fatty acids altered the changes in mRNA levels of all the four enzymes of the kynurenine pathway that had been modulated by IL-1 β exposure. EPA, DHA, sertraline and venlafaxine decreased the upregulation of IDO expression by 36%, 81%, 44% and 38% respectively, when compared with IL-1 β alone (Fig. 5a). Co-treatment with DHA, sertraline and venlafaxine decreased the upregulation of KMO caused by IL-1 β by 69%, 72% and 44%, respectively (Fig. 5b). Surprisingly, co-incubation with DHA caused a further 108% upregulation of KYNU when compared with IL-1 β alone (Fig. 5c). Finally, only sertraline and venlafaxine

reduced the increased in ACMSD caused by the cytokine, with changes of 44% and 71%, respectively (Fig. 5d).

4. Discussion

Our results show that EPA, DHA, sertraline and venlafaxine counteract the detrimental effects of IL-1 β on our *in vitro* model of human hippocampal neurogenesis.

EPA, DHA, sertraline and venlafaxine lessened the reduction in DCX-positive neuroblasts and MAP2-positive neurons caused by IL-1 β . This evidence that monoaminergic antidepressants and ω -3 fatty acids can reverse the anti-neurogenic effects of a pro-inflammatory cytokine on human hippocampal cells is of relevance given the importance attributed to immune dysfunctions in the pathogenesis of depression. Direct exposure to IL-1 β or to its inducers, including LPS and IFN- α , has been shown to suppress hippocampal neurogenesis, along with the development of depressive-like symptoms in animal models (Goshen et al., 2008; Koo and Duman, 2008; Ekdahl et al., 2009; Monje et al., 2003; Kaneko et al., 2006). Nonetheless, despite these observations and work demonstrating that IL-1 β is integral to the anhedonic and anti-neurogenic effects of stress (Koo and Duman, 2008), most evaluations of the protective effect of antidepressants in both animal and cellular experiments have modeled depression using stress paradigms or glucocorticoid exposure to simulate stressors. As such, chronic treatment with venlafaxine reversed the decrease in neurogenesis caused by chronic mild stress in mice (Zhang et al., 2015). Similarly, the suppressive effects of corticosterone on adult rat hippocampal cell proliferation were restored by paroxetine, another SSRI (Qiu et al., 2007). Additionally, we have shown that sertraline was able to restore the reductions of human hippocampal neurogenesis caused by *in vitro* treatment with dexamethasone in the same cellular model used for the present paper (Anacker et al., 2011). Only a few studies have used inflammatory models, aiming to efficiently mimic stress-related effects. Pre-treatment of human-like neuronal cells with trazodone, a tetracyclic antidepressant, completely reversed the decrease in cell viability caused by LPS and TNF- α (Daniele et al., 2015). Likewise, sertraline protected fetal rat hippocampal stem cells against LPS-induced cellular damage (Peng et al., 2012).

Inflammation has been thought to affect neurogenesis via several potential pathways: by activating the hypothalamic pituitary axis (HPA) axis to release cortisol and in turn reducing neurogene-




Fig. 4. Antidepressants and ω -3 fatty acids modulate changes of kynurenine pathway metabolites caused by IL-1 β treatment. Co-treatment of cells with IL-1 β and ω -3 fatty acids partially abrogated the increase in kynurenine caused by the cytokine (a). Co-treatment with sertraline fully reverted the decrease in levels of ANA caused by IL-1 β (b). Co-incubation with DHA and sertraline partially reverted the increase in QUIN levels caused by IL-1 β (c). No significant changes in PIC were observed by any co-incubation (d); * $p < 0.05$, ** $p < 0.001$.

sis; by affecting glial cell function, regulating the production of neurotrophic factors; and by direct effects on neural progenitor cells (Song and Wang, 2011). The present *in vitro* study excludes systemic effects such as changes to the HPA axis, so it allows for the examination of direct effects upon progenitor cells. DHA and sertraline reversed the effect of IL-1 β on reducing neurogenesis while also decreasing the production of quinolinic acid. At the same time, they caused a down-regulation of KMO mRNA expression. Hence, the ability of these compounds to block stimulation of quinolinic acid production or release induced by IL-1 β may be the mechanism by which they block the cytokine-induced reduction in neurogenesis. Surprisingly, ACMSD, known to enhance the production of the neuroprotective picolinic acid, was significantly upregulated by IL-1 β but also by co-treatment with IL-1 β and EPA or DHA, perhaps suggesting the inductions of a counteracting mechanisms against the increase in quinolinic acid previously reported. However, the levels of the picolinic acid were not affected in our experiments. This might suggest that either the increase in ACMSD was not strong enough to induce an effect on the metabolite level, or that perhaps a change in ACMSD gene expression requires a longer time to modulate levels of picolinic acid. While activation of the kynurenine pathway can lead to a reduction in serotonin, the behavioral and neural effects observed upon inflammatory insults are believed to be primarily dependent on the formation of neurotoxic metabolites (O'Connor et al., 2009; Dantzer et al.,

2011). In particular, quinolinic acid can increase neurotoxicity by various pathways, most notably by over-activation of NMDA receptors and increasing cytosolic calcium ions, leading to mitochondrial dysfunction, cytochrome *c* release, ATP exhaustion, free radical formation and oxidative damage (Perez-De La Cruz et al., 2012; Guillemin, 2012). Studies support the role of antidepressants acting on kynurenine metabolites, though they present a mixed picture. Evaluations in humans have shown that the reduced kynurenine acid/kynurenine neuroprotective ratio observed in depressed patients increased significantly following successful treatment with monoaminergic antidepressants (Myint et al., 2007). Similarly, treatment of patients with escitalopram showed a reduction of their initially elevated 3-hydroxykynurenine and quinolinic acid levels (Halaris et al., 2015). However, no effect of monoaminergic antidepressants was observed on levels of metabolites secreted by blood from depressed patients stimulated *in vitro* by LPS (Krause et al., 2012). Likewise, evaluations in animals have shown that imipramine reduced the initially elevated kynurenine/tryptophan ratio in the hippocampus of rats exposed to LPS or chronic mild stress (Elgarf et al., 2014), but caused no metabolite changes in a related study where the depression-like behavior was induced by IFN- α (Fischer et al., 2015). The regulation of kynurenine enzymes has also been measured: fluoxetine had no effect on KMO mRNA levels when used to prevent the development of depressive-like behavior in a mouse model of cancer related fati-




Fig. 5. Antidepressants and ω -3 fatty acids modulate IL-1 β -induced changes to the expression of enzymes of the kynurenine pathway. Co-treatment of cells with IL-1 β and ω -3 fatty acids or antidepressants decreased the expression of IDO (a). Co-treatment with DHA, sertraline or venlafaxine decreased expression of KMO (b). Co-treatment with DHA upregulated KMO production (c). Co-incubation with sertraline or venlafaxine decreased the upregulation of ACMSD (d); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

gue (Norden et al., 2015), while agomelatine, a melatonergic antidepressant, reversed the LPS-dependent increase in KMO expression in the hippocampus of rats (Molteni et al., 2013). Though conflicting, these studies are encouraging and support the notion that metabolites and enzymes of the kynurenine pathway are potential significant targets for therapeutic intervention. Interestingly, a recent study demonstrated that both EPA and DHA could increase neuronal differentiation in rat neural progenitor cells, but seemed to do so by alternative mechanisms, acting on different regulating genes to induce cell cycle arrest and neuronal differentiation (Katakura et al., 2013). Further investigation of the effect of these fatty acids, as well as a study of the action of sertraline and venlafaxine on cell cycle genes would be instructive.

Our findings are particularly relevant in understanding some of the clinical evidence related to the use of fish oils, and support previous evidence showing a positive effect of treatment with fatty acids on neuroinflammation, particularly in the context of depression (Mingam et al., 2008; Delpech et al., 2015; Labrousse et al., 2012; Larrieu et al., 2014; Bazinet and Laye, 2014; Orr et al., 2013; Lotrich et al., 2013; Kiecolt-Glaser et al., 2007; Freeman and Rapaport, 2011; Mocking et al., 2017). Recent meta-analyses suggested that EPA has better antidepressant effects than DHA in combination with antidepressant medications (Mocking et al., 2016; Grosso et al., 2014). Indeed, the clinical trials recruiting depressed patients taking antidepressant agents as a combination

showed that either DHA is ineffective (Grenyer et al., 2007; Silvers et al., 2005) or EPA is more effective than DHA (Mozaffari-Khosravi et al., 2013). In addition, it has been reported that the combined effect of higher dose of EPA with lower dose of DHA was effective in reducing depressive symptoms in depression (Rondanelli et al., 2010; Rizzo et al., 2012) whereas the opposite combination (lower EPA and higher DHA) was ineffective (Silvers et al., 2005; Su et al., 2003; Grenyer et al., 2007), perhaps suggesting that the dosage, more than the type of treatment, might influence patients' response. Our present work shows that treatment with either DHA or EPA has a positive effect on neurogenesis, with DHA also decreasing the production of the neurotoxic quinolinic acid. In contrast, in our previous study we demonstrated an overall anti-inflammatory pattern for EPA, but a mostly pro-inflammatory one for DHA upon co-treatment with IL-1 β . Interestingly, both those effects were associated with a corresponding decrease in the NF- κ B activation caused by IL-1 β alone (Horowitz et al., 2015). In addition, we have also seen that EPA, but not DHA, can revert the damage caused by oxidative stress in our human hippocampal cell model. We have no indication about the overall status of the patients evaluated in clinical studies (including neurogenesis, oxidative stress, inflammation or regulation in the kynurenine pathway), so a possible overlap or the prominence of any of the possible dysregulations cannot be excluded. Considering that all clinical studies involved supplementation, it is also pos-

sible that EPA and DHA have differential interactions with various antidepressant drugs. Studies with individual combinations would therefore be of interest.

5. Conclusions

In summary, our observations indicate that venlafaxine, sertraline, EPA and DHA show protective effects against the detrimental influence of IL-1 β on human hippocampal neurogenesis. We also find that one potential explanation for these effects is that these compounds may be involved in regulating IL-1 β signaling by modulation of the kynurenine pathway, as DHA and sertraline reduce levels of the neurotoxic quinolinic acid increased by the cytokine, with a similar trend observed for EPA and venlafaxine. An increased understanding of the molecular mechanisms underlying these effects may allow for a more effective future personalization of antidepressant use.

Conflict of interest

Professor Pariente and Dr. Zunszain have received research funding from Johnson & Johnson as part of a program of research on depression and inflammation, and research funding from the Medical Research Council (UK) and the Wellcome Trust for research on depression and inflammation as part of two large consortia that also include Johnson & Johnson, GSK and Lundbeck. The work presented in this paper is unrelated to this funding. All other authors declare no conflict of interest.

Acknowledgments

This study was supported in part by a Joint Research Project (102-2911-I-039-501) between the Ministry of Science and Technology (Taiwan) and the Royal Society (UK), by the Medical Research Council (UK) grant 'AMBROSIAC – A Menu for Brain Responses Opposing Stress-Induced Alterations in Cognition' (MR/N029488/1), and by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health.

References

- Boldrini, M. et al., 2009. Antidepressants increase neural progenitor cells in the human hippocampus. *Neuropsychopharmacology* 34 (11), 2376–2389.
- Kempermann, G., Krebs, J., Fabel, K., 2008. The contribution of failing adult hippocampal neurogenesis to psychiatric disorders. *Curr. Opin. Psychiatry* 21 (3), 290–295.
- Hill, A.S., Sahay, A., Hen, R., 2015. Increasing adult hippocampal neurogenesis is sufficient to reduce anxiety and depression-like behaviors. *Neuropsychopharmacology* 40 (10), 2368–2378.
- Warner-Schmidt, J.L., Duman, R.S., 2006. Hippocampal neurogenesis: opposing effects of stress and antidepressant treatment. *Hippocampus* 16 (3), 239–249.
- Sahay, A., Hen, R., 2007. Adult hippocampal neurogenesis in depression. *Nat. Neurosci.* 10 (9), 1110–1115.
- Malberg, J.E., Blendy, J.A., 2005. Antidepressant action: to the nucleus and beyond. *Trends Pharmacol. Sci.* 26 (12), 631–638.
- Goshen, I. et al., 2008. Brain interleukin-1 mediates chronic stress-induced depression in mice via adrenocortical activation and hippocampal neurogenesis suppression. *Mol. Psychiatry* 13 (7), 717–728.
- Koo, J.W., Duman, R.S., 2008. IL-1 β is an essential mediator of the antineurogenic and anhedonic effects of stress. *Proc. Natl. Acad. Sci. U.S.A.* 105 (2), 751–756.
- Kuzumaki, N. et al., 2008. Enhanced IL-1 β production in response to the activation of hippocampal glial cells impairs neurogenesis in aged mice. *Synapse* 64 (9), 721–728.
- Capuron, L., Miller, A.H., 2004. Cytokines and psychopathology: lessons from interferon-alpha. *Biol. Psychiatry* 56 (11), 819–824.
- Howren, M.B., Lamkin, D.M., Suls, J., 2009. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom. Med.* 71 (2), 171–186.
- Maes, M. et al., 2009. The inflammatory & neurodegenerative (I&ND) hypothesis of depression: leads for future research and new drug developments in depression. *Metab. Brain Dis.* 24 (1), 27–53.
- Miller, A.H., Maletic, V., Raison, C.L., 2009. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol. Psychiatry* 65 (9), 732–741.
- Mossner, R. et al., 2007. Consensus paper of the WFSBP Task Force on Biological Markers: biological markers in depression. *World J. Biol. Psychiatry* 8 (3), 141–174.
- Raison, C.L., Capuron, L., Miller, A.H., 2006. Cytokines sing the blues: inflammation and the pathogenesis of depression. *Trends Immunol.* 27 (1), 24–31.
- Schiepers, O.J., Wichers, M.C., Maes, M., 2005. Cytokines and major depression. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 29 (2), 201–217.
- Yirmiya, R., Goshen, I., 2011. Immune modulation of learning, memory, neural plasticity and neurogenesis. *Brain Behav. Immun.* 25 (2), 181–213.
- Schwarcz, R. et al., 2012. Kynurenines in the mammalian brain: when physiology meets pathology. *Nat. Rev. Neurosci.* 13 (7), 465–477.
- Fischer, C.W. et al., 2015. Interferon-alpha treatment induces depression-like behaviour accompanied by elevated hippocampal quinolinic acid levels in rats. *Behav. Brain Res.* 293, 166–172.
- Raison, C.L. et al., 2010. CSF concentrations of brain tryptophan and kynurenines during immune stimulation with IFN-alpha: relationship to CNS immune responses and depression. *Mol. Psychiatry* 15 (4), 393–403.
- Erhardt, S. et al., 2013. Connecting inflammation with glutamate agonism in suicidality. *Neuropsychopharmacology* 38 (5), 743–752.
- Brundin, L. et al., 2016. An enzyme in the kynurenine pathway that governs vulnerability to suicidal behavior by regulating excitotoxicity and neuroinflammation. *Transl. Psychiatry* 6 (8), e865.
- Capuron, L. et al., 2003. Interferon-alpha-induced changes in tryptophan metabolism: relationship to depression and paroxetine treatment. *Biol. Psychiatry* 54 (9), 906–914.
- Guillemin, G.J. et al., 2005. Indoleamine 2,3-dioxygenase and quinolinic acid immunoreactivity in Alzheimer's disease hippocampus. *Neuropathol. Appl. Neurobiol.* 31 (4), 395–404.
- Meier, T.B. et al., 2016. Relationship between neurotoxic kynurenine metabolites and reductions in right medial prefrontal cortical thickness in major depressive disorder. *Brain Behav. Immun.* 53, 39–48.
- Savitz, J. et al., 2015. Reduction of kynurenic acid to quinolinic acid ratio in both the depressed and remitted phases of major depressive disorder. *Brain Behav. Immun.* 46, 55–59.
- O'Connor, J.C. et al., 2009. Lipopolysaccharide-induced depressive-like behavior is mediated by indoleamine 2,3-dioxygenase activation in mice. *Mol. Psychiatry* 14 (5), 511–522.
- Dobos, N. et al., 2012. The role of indoleamine 2,3-dioxygenase in a mouse model of neuroinflammation-induced depression. *J. Alzheimers Dis.* 28 (4), 905–915.
- Zwilling, D. et al., 2011. Kynurenine 3-monooxygenase inhibition in blood ameliorates neurodegeneration. *Cell* 145 (6), 863–874.
- Heisler, J.M., O'Connor, J.C., 2015. Indoleamine 2,3-dioxygenase-dependent neurotoxic kynurenine metabolism mediates inflammation-induced deficit in recognition memory. *Brain Behav. Immun.* 50, 115–124.
- Parrott, J.M. et al., 2016. Neurotoxic kynurenine metabolism is increased in the dorsal hippocampus and drives distinct depressive behaviors during inflammation. *Transl. Psychiatry* 6 (10), e918.
- Zunszain, P.A. et al., 2012. Interleukin-1 β : a new regulator of the kynurenine pathway affecting human hippocampal neurogenesis. *Neuropsychopharmacology* 37 (4), 939–949.
- Anacker, C. et al., 2011. Antidepressants increase human hippocampal neurogenesis by activating the glucocorticoid receptor. *Mol. Psychiatry*.
- Hannestad, J., DellaGioia, N., Bloch, M., 2011. The effect of antidepressant medication treatment on serum levels of inflammatory cytokines: a meta-analysis. *Neuropsychopharmacology* 36 (12), 2452–2459.
- Su, K.P. et al., 2014. Omega-3 fatty acids in the prevention of interferon-alpha-induced depression: results from a randomized, controlled trial. *Biol. Psychiatry* 76 (7), 559–566.
- Su, K.P., 2012. Inflammation in psychopathology of depression: clinical, biological, and therapeutic implications. *BioMedicine* 2 (2), 68–74.
- Lin, P.Y. et al., 2012. Are omega-3 fatty acids antidepressants or just mood-improving agents? The effect depends upon diagnosis, supplement preparation, and severity of depression. *Mol. Psychiatry* 17 (12), 1161–1163.
- Lu, D.Y. et al., 2010. Docosahexaenoic acid suppresses neuroinflammatory responses and induces heme oxygenase-1 expression in BV-2 microglia: implications of antidepressant effects for omega-3 fatty acids. *Neuropsychopharmacology* 35 (11), 2238–2248.
- Lin, P.Y., Huang, S.Y., Su, K.P., 2010. A meta-analytic review of polyunsaturated fatty acid compositions in patients with depression. *Biol. Psychiatry* 68 (2), 140–147.
- Su, K.P., 2009. Biological mechanism of antidepressant effect of omega-3 fatty acids: how does fish oil act as a 'mind-body interface'? *Neurosignals* 17 (2), 144–152.
- Horowitz, M.A. et al., 2015. Antidepressant compounds can be both pro- and anti-inflammatory in human hippocampal cells. *Int. J. Neuropsychopharmacol.* 18 (3).
- Anacker, C. et al., 2011. Antidepressants increase human hippocampal neurogenesis by activating the glucocorticoid receptor. *Mol. Psychiatry* 16 (7), 738–750.

- Bustin, S.A. et al., 2005. Quantitative real-time RT-PCR—a perspective. *J. Mol. Endocrinol.* 34 (3), 597–601.
- Ekdahl, C.T., Kokaia, Z., Lindvall, O., 2009. Brain inflammation and adult neurogenesis: the dual role of microglia. *Neuroscience* 158 (3), 1021–1029.
- Monje, M.L., Toda, H., Palmer, T.D., 2003. Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302 (5651), 1760–1765.
- Kaneko, N. et al., 2006. Suppression of cell proliferation by interferon- α through interleukin-1 production in adult rat dentate gyrus. *Neuropsychopharmacology* 31 (12), 2619–2626.
- Zhang, X. et al., 2015. Venlafaxine increases cell proliferation and regulates DISC1, PDE4B and NMDA receptor 2B expression in the hippocampus in chronic mild stress mice. *Eur. J. Pharmacol.* 755, 58–65.
- Qiu, G. et al., 2007. Modulation of the suppressive effect of corticosterone on adult rat hippocampal cell proliferation by paroxetine. *Neurosci. Bull.* 23 (3), 131–136.
- Daniele, S. et al., 2015. Trazodone treatment protects neuronal-like cells from inflammatory insult by inhibiting NF- κ B, p38 and JNK. *Cell. Signal.* 27 (8), 1609–1629.
- Peng, Z.W. et al., 2012. Sertraline promotes hippocampus-derived neural stem cells differentiating into neurons but not glia and attenuates LPS-induced cellular damage. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 36 (1), 183–188.
- Song, C., Wang, H., 2011. Cytokines mediated inflammation and decreased neurogenesis in animal models of depression. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 35 (3), 760–768.
- Dantzer, R. et al., 2011. Inflammation-associated depression: from serotonin to kynurenine. *Psychoneuroendocrinology* 36 (3), 426–436.
- Perez-De La Cruz, V., Carrillo-Mora, P., Santamaria, A., 2012. Quinolinic acid, an endogenous molecule combining excitotoxicity, oxidative stress and other toxic mechanisms. *Int. J. Tryptophan Res.* 5, 1–8.
- Guillemin, G.J., 2012. Quinolinic acid, the inescapable neurotoxin. *FEBS J.* 279 (8), 1356–1365.
- Myint, A.M. et al., 2007. Kynurenine pathway in major depression: evidence of impaired neuroprotection. *J. Affect. Disord.* 98 (1–2), 143–151.
- Halaris, A. et al., 2015. Does escitalopram reduce neurotoxicity in major depression? *J. Psychiatr. Res.* 66–67, 118–126.
- Krause, D.L. et al., 2012. Effects of antidepressants and cyclooxygenase-2 inhibitor on cytokines and kynurenines in stimulated in vitro blood culture from depressed patients. *Inflammopharmacology* 20 (3), 169–176.
- Elgarf, A.S. et al., 2014. Lipopolysaccharide repeated challenge followed by chronic mild stress protocol introduces a combined model of depression in rats: reversibility by imipramine and pentoxifylline. *Pharmacol. Biochem. Behav.* 126, 152–162.
- Norden, D.M. et al., 2015. Fluoxetine prevents the development of depressive-like behavior in a mouse model of cancer related fatigue. *Physiol. Behav.* 140, 230–235.
- Molteni, R. et al., 2013. Modulation of the inflammatory response in rats chronically treated with the antidepressant agomelatine. *Eur. Neuropsychopharmacol.* 23 (11), 1645–1655.
- Katakura, M. et al., 2013. Omega-3 polyunsaturated Fatty acids enhance neuronal differentiation in cultured rat neural stem cells. *Stem Cells Int.* 2013, 490476.
- Mingam, R. et al., 2008. Uncoupling of interleukin-6 from its signalling pathway by dietary n-3-polyunsaturated fatty acid deprivation alters sickness behaviour in mice. *Eur. J. Neurosci.* 28 (9), 1877–1886.
- Delpech, J.C. et al., 2015. Dietary n-3 PUFAs deficiency increases vulnerability to inflammation-induced spatial memory impairment. *Neuropsychopharmacology* 40 (12), 2774–2787.
- Labrousse, V.F. et al., 2012. Short-term long chain omega3 diet protects from neuroinflammatory processes and memory impairment in aged mice. *PLoS ONE* 7 (5), e36861.
- Laureau, T. et al., 2014. Nutritional omega-3 modulates neuronal morphology in the prefrontal cortex along with depression-related behaviour through corticosterone secretion. *Transl. Psychiatry* 4, e437.
- Bazinet, R.P., Laye, S., 2014. Polyunsaturated fatty acids and their metabolites in brain function and disease. *Nat. Rev. Neurosci.* 15 (12), 771–785.
- Orr, S.K., Trepanier, M.O., Bazinet, R.P., 2013. n-3 polyunsaturated fatty acids in animal models with neuroinflammation. *Prostaglandins Leukot. Essent. Fatty Acids* 88 (1), 97–103.
- Lotrich, F.E., Sears, B., McNamara, R.K., 2013. Elevated ratio of arachidonic acid to long-chain omega-3 fatty acids predicts depression development following interferon- α treatment: relationship with interleukin-6. *Brain Behav. Immun.* 31, 48–53.
- Kiecolt-Glaser, J.K. et al., 2007. Depressive symptoms, omega-6:omega-3 fatty acids, and inflammation in older adults. *Psychosom. Med.* 69 (3), 217–224.
- Freeman, M.P., Rapaport, M.H., 2011. Omega-3 fatty acids and depression: from cellular mechanisms to clinical care. *J. Clin. Psychiatry* 72 (2), 258–259.
- Mocking, R.J. et al., 2017. Biological profiling of prospective antidepressant response in major depressive disorder: associations with (neuro)inflammation, fatty acid metabolism, and amygdala-reactivity. *Psychoneuroendocrinology* 79, 84–92.
- Mocking, R.J. et al., 2016. Meta-analysis and meta-regression of omega-3 polyunsaturated fatty acid supplementation for major depressive disorder. *Transl. Psychiatry* 6, e756.
- Grosso, G. et al., 2014. Role of omega-3 fatty acids in the treatment of depressive disorders: a comprehensive meta-analysis of randomized clinical trials. *PLoS ONE* 9 (5), e96905.
- Grenyer, B.F. et al., 2007. Fish oil supplementation in the treatment of major depression: a randomised double-blind placebo-controlled trial. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 31 (7), 1393–1396.
- Silvers, K.M. et al., 2005. Randomised double-blind placebo-controlled trial of fish oil in the treatment of depression. *Prostaglandins Leukot. Essent. Fatty Acids* 72 (3), 211–218.
- Mozaffari-Khosravi, H. et al., 2013. Eicosapentaenoic acid versus docosahexaenoic acid in mild-to-moderate depression: a randomized, double-blind, placebo-controlled trial. *Eur. Neuropsychopharmacol.* 23 (7), 636–644.
- Rondanelli, M. et al., 2010. Effect of omega-3 fatty acids supplementation on depressive symptoms and on health-related quality of life in the treatment of elderly women with depression: a double-blind, placebo-controlled, randomized clinical trial. *J. Am. Coll. Nutr.* 29 (1), 55–64.
- Rizzo, A.M. et al., 2012. Comparison between the AA/EPA ratio in depressed and non depressed elderly females: omega-3 fatty acid supplementation correlates with improved symptoms but does not change immunological parameters. *Nutr. J.* 11, 82.
- Silvers, K.M. et al., 2005. Randomised double-blind placebo-controlled trial of fish oil in the treatment of depression. *Prostaglandins Leukot. Essent. Fatty Acids* 72 (3), 211–218.
- Su, K.P. et al., 2003. Omega-3 fatty acids in major depressive disorder. A preliminary double-blind, placebo-controlled trial. *Eur. Neuropsychopharmacol.* 13 (4), 267–271.
- Grenyer, B.F. et al., 2007. Fish oil supplementation in the treatment of major depression: a randomised double-blind placebo-controlled trial. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 31 (7), 1393–1396.